arXiv:astro-ph/0406033v1 1 Jun 2004

THE $Z > 4$ QUASAR POPULATION OBSERVED BY CHANDRA AND XMM-NEWTON

C. VIGNALI*
INAF–Osservatorio Astronomico di Bologna,
Via Ranzani, 1 – I-40127 Bologna, Italy
E-mail: cristian.vignali@bo.astro.it

W.N. BRANDT AND D.P. SCHNEIDER
Department of Astronomy & Astrophysics,
The Pennsylvania State University,
525 Davey Lab, University Park, PA 16802, USA
E-mail: niel,dps@astro.psu.edu

The current status of our *Chandra* and XMM-*Newton* project on high-redshift ($z \gtrsim 4$) quasars is briefly reviewed. We report the main results obtained in the last few years for the X-ray detected quasars, along with a few ($\approx 10\%$) intriguing cases where no detection has been obtained with *Chandra* snapshot observations.

1. Introduction

Quasars at $z \gtrsim 4$ provide direct information on the first massive structures to form in the Universe. The last few years have evidenced incredible progress in the study of the high-redshift Universe, largely thanks to ground-based optical surveys. In particular, the Sloan Digital Sky Survey (SDSS[1]) has discovered large numbers of high-redshift quasars, increasing the number of known quasars at $z \gtrsim 4$ to > 500,[2] and many more are expected to be discovered before the survey ends (see [3] for the recent advances provided by the SDSS). Many of the high-redshift quasars discovered by recent optical surveys are suitable for follow-up X-ray investigations. Here we review X-ray studies of the highest redshift quasars, focusing on recent advances enabled largely by the imaging and spectroscopic capabilities of *Chandra* and XMM-*Newton*.

---

*Work partially supported by the Italian Space Agency under the contract ASI I/R/057/02 (CV) and by the Chandra X-ray Center grant GO2-3134X (WNB).



Figure 1. Observed-frame, Galactic absorption-corrected 0.5–2 keV flux versus $AB_{1450(1+z)}$ magnitude for $z \gtrsim 4$ AGN found in optical (triangles) and X-ray (squares) surveys. The rapid increase in the number of X-ray detected AGN at $z \gtrsim 4$ is evident when the two panels (i.e., the situation in year 2000 vs. the current one) are compared. Most of the upper limits have been obtained from archival *ROSAT* observations.

2. X-ray detections at $z \gtrsim 4$

Our knowledge of the X-ray properties of $z \gtrsim 4$ quasars has advanced rapidly over the last few years. Since the pioneering work based on the systematic analysis of archival *ROSAT* data[4], where the number of X-ray detected quasars at $z > 4$ doubled, the progress made in this field has been substantial (see Fig. 1). At present, more than 90 quasars at $z \gtrsim 4$ have X-ray detections;[a] while in most cases *Chandra* snapshot (typically 4–6 ks) observations have allowed derivation of basic X-ray information (X-ray fluxes and luminosities),[5,6,7,8,9,10,11,12,13] for a few X-ray luminous objects medium-quality X-ray spectra have also been obtained[14,15] thanks to the large collecting area of XMM-*Newton*. From a general perspective, the most important results from X-ray analyses of high-redshift quasars are the following:

◇ The X-ray properties of $z \gtrsim 4$ quasars are similar to those of local quasars. Joint X-ray spectral fitting of optically luminous quasars at $z \gtrsim 4$ selected from the Palomar Digital Sky Survey (DPOSS[16]) and the SDSS has shown[10,11] that quasar X-ray continuum shapes do not show evidence for significant spectral evolution over cosmic time. This has found further support recently via joint spectral fitting of a sample of 46 radio-quiet quasars (RQQs) with *Chandra* detections in the redshift range 4.0–6.3 (≈ 750 source counts; $\Gamma = 1.9 \pm 0.1$)[17] and by direct X-ray spectral analyses of a few objects observed by XMM-*Newton*.[14,15] No evidence for widespread X-ray absorption has been found, although some quasars (see discussion below) are likely to be obscured.

---

[a] See http://www.astro.psu.edu/users/niel/papers/highz-xray-detected.dat for a regularly updated listing of X-ray detections and sensitive upper limits at $z \gtrsim 4$.

◇ The optical-to-X-ray properties of high-redshift quasars and those of local quasars have been found similar, once luminosity effects and selection biases are taken into account properly.[18] Overall, the emerging picture is that the small-scale X-ray emission regions of quasars appear relatively insensitive to large-scale environmental differences at $z \approx 6$.

◇ Moderately deep *Chandra* observations and the ultra-deep (2 Ms) survey in the *Chandra* Deep Field-North (CDF-N)[19] have allowed X-ray studies of moderate-luminosity AGN at $z > 4$; this population of AGN is more numerous and thus more representative than the rare, highly luminous quasars.[20]

◇ Similarly to the results obtained for the RQQs, neither the X-ray spectral slope ($\Gamma = 1.65 \pm 0.15$) nor the jet emission of the radio-loud quasar population seems to evolve with cosmic time.[21] No evidence for significant X-ray brightening ascribed to inverse Compton scattering of energetic electrons with Cosmic Microwave Background photons has been revealed by snapshot observations with *Chandra*.[21]

3. X-ray non-detections: space oddities at high redshift?

At present, the fraction of X-ray upper limits among the sensitive *Chandra* and XMM-*Newton* observations of high-redshift quasars is low ($\approx 10\%$). About half of the X-ray undetected quasars are broad absorption-line quasars (BALQSOs), which are known to be absorbed in the X-ray band;[22] one quasar shows narrow absorption features from N V and Lyα at approximately the source redshift.[11] For most of these objects the lack of X-ray detections at the flux limits reached by *Chandra* with snapshot observations is likely due to absorption by column densities larger[6,11] than $\approx 10^{23}$ cm^{-2} (assuming reasonable values for the optical-to-X-ray spectral energy distribution and X-ray photon index). It is possible that the remaining X-ray undetected high-redshift quasars are characterized by absorption features not recognized in low-resolution optical spectra. Recently, it has been shown that near-infrared spectroscopy can be highly effective[23] in detecting absorption features in high-redshift objects previously classified as "normal" quasars. Clearly, minority AGN populations at $z > 4$ need to be investigated better in the X-ray regime. These include BALQSOs (whose number is still too limited to derive statistically reliable average properties) and the population of quasars lacking emission lines.[24,6] Follow-up observations of quasars without optical emission lines with *Chandra* are on-going and the results will be presented in a forthcoming paper.[25]

References

1. D.G. York *et al.*, *AJ* **120**, 1579 (2000).
2. X. Fan, private communication (2004).
3. K. Abazajian *et al.*, *AJ*, in press, astro-ph/0403325 (2004).
4. S. Kaspi, W.N. Brandt and D.P. Schneider, *AJ* **119**, 2031 (2000; KBS00).
5. W.N. Brandt *et al.*, *AJ* **121**, 591 (2001).
6. C. Vignali, W.N. Brandt, X. Fan, J.E. Gunn, S. Kaspi, D.P. Schneider and M.A. Strauss, *AJ* **122**, 2143 (2001).
7. W.N. Brandt *et al.*, *ApJ* **569**, L5 (2002).
8. J.D. Silverman *et al.*, *ApJ* **569**, L1 (2002).
9. J. Bechtold *et al.*, *ApJ* **588**, 119 (2003).
10. C. Vignali, W.N. Brandt, D.P. Schneider, G.P. Garmire and S. Kaspi, *AJ* **125**, 418 (2003).
11. C. Vignali *et al.*, *AJ* **125**, 2876 (2003).
12. F.J. Castander, E. Treister, T.J. Maccarone, P.S. Coppi, J. Maza, S.E. Zepf and R. Guzman, *AJ* **125**, 1689 (2003).
13. E. Treister, F.J. Castander, T.J. Maccarone, D. Herrera, E. Gawiser, J. Maza and P.S. Coppi, *ApJ* **603**, 36 (2004).
14. E. Ferrero and W. Brinkmann, *A&A* **402**, 465 (2003).
15. D. Grupe, S. Mathur, B.J. Wilkes and M. Elvis, *AJ* **127**, 1 (2004).
16. S.G. Djorgovski *et al.*, in Proceedings of "Wide Field Surveys in Cosmology", 89 (1998).
17. C. Vignali, W.N. Brandt and D.P. Schneider, in Proceedings of "AGN Physics with the Sloan Digital Sky Survey", in press, astro-ph/0310659 (2004).
18. C. Vignali, W.N. Brandt and D.P. Schneider, *AJ* **125**, 433 (2003).
19. D.M. Alexander *et al.*, *AJ* **126**, 539 (2003).
20. C. Vignali, F.E. Bauer, D.M. Alexander, W.N. Brandt, A.E. Hornschemeier, D.P. Schneider and G.P. Garmire, *ApJ* **580**, L105 (2002).
21. L.C. Bassett, W.N. Brandt, D.P. Schneider, C. Vignali and G.P. Garmire, *AJ*, in press, astro-ph/0404543 (2004).
22. W.N. Brandt, A. Laor and B.J. Wills, *ApJ* **528**, 637 (2000).
23. R. Maiolino, E. Oliva, F. Ghinassi, M. Pedani, F. Mannucci, R. Mujica and Y. Juarez, *A&A*, in press, astro-ph/0312402 (2004).
24. X. Fan *et al.*, *ApJ* **526**, L57 (1999).
25. W.N. Brandt *et al.*, in preparation.